

Exercise 7.7

Question 1:

$$\sqrt{4-x^2}$$

Answer

$$\text{Let } I = \int \sqrt{4-x^2} dx = \int \sqrt{(2)^2 - (x)^2} dx$$

$$\text{It is known that, } \int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + C$$

$$\begin{aligned} \therefore I &= \frac{x}{2} \sqrt{4-x^2} + \frac{4}{2} \sin^{-1} \frac{x}{2} + C \\ &= \frac{x}{2} \sqrt{4-x^2} + 2 \sin^{-1} \frac{x}{2} + C \end{aligned}$$

Question 2:

$$\sqrt{1-4x^2}$$

Answer

$$\text{Let } I = \int \sqrt{1-4x^2} dx = \int \sqrt{(1)^2 - (2x)^2} dx$$

$$\text{Let } 2x = t \Rightarrow 2 dx = dt$$

$$\therefore I = \frac{1}{2} \int \sqrt{(1)^2 - (t)^2} dt$$

$$\text{It is known that, } \int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + C$$

$$\Rightarrow I = \frac{1}{2} \left[\frac{t}{2} \sqrt{1-t^2} + \frac{1}{2} \sin^{-1} t \right] + C$$

$$= \frac{t}{4} \sqrt{1-t^2} + \frac{1}{4} \sin^{-1} t + C$$

$$= \frac{2x}{4} \sqrt{1-4x^2} + \frac{1}{4} \sin^{-1} 2x + C$$

$$= \frac{x}{2} \sqrt{1-4x^2} + \frac{1}{4} \sin^{-1} 2x + C$$

Question 3:

$$\sqrt{x^2 + 4x + 6}$$

Answer

$$\begin{aligned}\text{Let } I &= \int \sqrt{x^2 + 4x + 6} \, dx \\ &= \int \sqrt{x^2 + 4x + 4 + 2} \, dx \\ &= \int \sqrt{(x^2 + 4x + 4) + 2} \, dx \\ &= \int \sqrt{(x+2)^2 + (\sqrt{2})^2} \, dx\end{aligned}$$

$$\text{It is known that, } \int \sqrt{x^2 + a^2} \, dx = \frac{x}{2} \sqrt{x^2 + a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2 + a^2}| + C$$

$$\begin{aligned}\therefore I &= \frac{(x+2)}{2} \sqrt{x^2 + 4x + 6} + \frac{2}{2} \log |(x+2) + \sqrt{x^2 + 4x + 6}| + C \\ &= \frac{(x+2)}{2} \sqrt{x^2 + 4x + 6} + \log |(x+2) + \sqrt{x^2 + 4x + 6}| + C\end{aligned}$$

Question 4:

$$\sqrt{x^2 + 4x + 1}$$

Answer

$$\begin{aligned}\text{Let } I &= \int \sqrt{x^2 + 4x + 1} \, dx \\ &= \int \sqrt{(x^2 + 4x + 4) - 3} \, dx \\ &= \int \sqrt{(x+2)^2 - (\sqrt{3})^2} \, dx\end{aligned}$$

$$\text{It is known that, } \int \sqrt{x^2 - a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log |x + \sqrt{x^2 - a^2}| + C$$

$$\therefore I = \frac{(x+2)}{2} \sqrt{x^2 + 4x + 1} - \frac{3}{2} \log |(x+2) + \sqrt{x^2 + 4x + 1}| + C$$

Question 5:

$$\sqrt{1-4x-x^2}$$

Answer

$$\begin{aligned}\text{Let } I &= \int \sqrt{1-4x-x^2} \, dx \\ &= \int \sqrt{1-(x^2+4x+4-4)} \, dx \\ &= \int \sqrt{1+4-(x+2)^2} \, dx \\ &= \int \sqrt{(\sqrt{5})^2 - (x+2)^2} \, dx\end{aligned}$$

It is known that, $\int \sqrt{a^2 - x^2} \, dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + C$

$$\therefore I = \frac{(x+2)}{2} \sqrt{1-4x-x^2} + \frac{5}{2} \sin^{-1} \left(\frac{x+2}{\sqrt{5}} \right) + C$$

Question 6:

$$\sqrt{x^2+4x-5}$$

Answer

$$\begin{aligned}\text{Let } I &= \int \sqrt{x^2+4x-5} \, dx \\ &= \int \sqrt{(x^2+4x+4)-9} \, dx \\ &= \int \sqrt{(x+2)^2 - (3)^2} \, dx\end{aligned}$$

It is known that, $\int \sqrt{x^2 - a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log |x + \sqrt{x^2 - a^2}| + C$

$$\therefore I = \frac{(x+2)}{2} \sqrt{x^2+4x-5} - \frac{9}{2} \log |(x+2) + \sqrt{x^2+4x-5}| + C$$

Question 7:

$$\sqrt{1+3x-x^2}$$

Answer

$$\begin{aligned}\text{Let } I &= \int \sqrt{1+3x-x^2} dx \\ &= \int \sqrt{1-\left(x^2-3x+\frac{9}{4}-\frac{9}{4}\right)} dx \\ &= \int \sqrt{\left(1+\frac{9}{4}\right)-\left(x-\frac{3}{2}\right)^2} dx \\ &= \int \sqrt{\left(\frac{\sqrt{13}}{2}\right)^2-\left(x-\frac{3}{2}\right)^2} dx\end{aligned}$$

It is known that, $\int \sqrt{a^2-x^2} dx = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + C$

$$\begin{aligned}\therefore I &= \frac{x-\frac{3}{2}}{2} \sqrt{1+3x-x^2} + \frac{13}{4 \times 2} \sin^{-1} \left(\frac{x-\frac{3}{2}}{\frac{\sqrt{13}}{2}} \right) + C \\ &= \frac{2x-3}{4} \sqrt{1+3x-x^2} + \frac{13}{8} \sin^{-1} \left(\frac{2x-3}{\sqrt{13}} \right) + C\end{aligned}$$

Question 8:

$$\sqrt{x^2+3x}$$

Answer

$$\begin{aligned}\text{Let } I &= \int \sqrt{x^2+3x} dx \\ &= \int \sqrt{x^2+3x+\frac{9}{4}-\frac{9}{4}} dx \\ &= \int \sqrt{\left(x+\frac{3}{2}\right)^2-\left(\frac{3}{2}\right)^2} dx\end{aligned}$$

It is known that, $\int \sqrt{x^2 - a^2} dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log \left| x + \sqrt{x^2 - a^2} \right| + C$

$$\begin{aligned}\therefore I &= \frac{\left(x + \frac{3}{2}\right)}{2} \sqrt{x^2 + 3x} - \frac{9}{2} \log \left| \left(x + \frac{3}{2}\right) + \sqrt{x^2 + 3x} \right| + C \\ &= \frac{(2x+3)}{4} \sqrt{x^2 + 3x} - \frac{9}{8} \log \left| \left(x + \frac{3}{2}\right) + \sqrt{x^2 + 3x} \right| + C\end{aligned}$$

Question 9:

$$\sqrt{1 + \frac{x^2}{9}}$$

Answer

$$\text{Let } I = \int \sqrt{1 + \frac{x^2}{9}} dx = \frac{1}{3} \int \sqrt{9 + x^2} dx = \frac{1}{3} \int \sqrt{(3)^2 + x^2} dx$$

It is known that, $\int \sqrt{x^2 + a^2} dx = \frac{x}{2} \sqrt{x^2 + a^2} + \frac{a^2}{2} \log \left| x + \sqrt{x^2 + a^2} \right| + C$

$$\begin{aligned}\therefore I &= \frac{1}{3} \left[\frac{x}{2} \sqrt{x^2 + 9} + \frac{9}{2} \log \left| x + \sqrt{x^2 + 9} \right| \right] + C \\ &= \frac{x}{6} \sqrt{x^2 + 9} + \frac{3}{2} \log \left| x + \sqrt{x^2 + 9} \right| + C\end{aligned}$$

Question 10:

$\int \sqrt{1+x^2} dx$ is equal to

A. $\frac{x}{2} \sqrt{1+x^2} + \frac{1}{2} \log \left| x + \sqrt{1+x^2} \right| + C$

B. $\frac{2}{3} (1+x^2)^{\frac{2}{3}} + C$

C. $\frac{2}{3} x (1+x^2)^{\frac{3}{2}} + C$

D. $\frac{x^2}{2}\sqrt{1+x^2} + \frac{1}{2}x^2 \log|x+\sqrt{1+x^2}| + C$

Answer

It is known that, $\int \sqrt{a^2+x^2} dx = \frac{x}{2}\sqrt{a^2+x^2} + \frac{a^2}{2} \log|x+\sqrt{x^2+a^2}| + C$

$\therefore \int \sqrt{1+x^2} dx = \frac{x}{2}\sqrt{1+x^2} + \frac{1}{2} \log|x+\sqrt{1+x^2}| + C$

Hence, the correct Answer is A.

Question 11:

$\int \sqrt{x^2-8x+7} dx$ is equal to

A. $\frac{1}{2}(x-4)\sqrt{x^2-8x+7} + 9 \log|x-4+\sqrt{x^2-8x+7}| + C$

B. $\frac{1}{2}(x+4)\sqrt{x^2-8x+7} + 9 \log|x+4+\sqrt{x^2-8x+7}| + C$

C. $\frac{1}{2}(x-4)\sqrt{x^2-8x+7} - 3\sqrt{2} \log|x-4+\sqrt{x^2-8x+7}| + C$

D. $\frac{1}{2}(x-4)\sqrt{x^2-8x+7} - \frac{9}{2} \log|x-4+\sqrt{x^2-8x+7}| + C$

Answer

Let $I = \int \sqrt{x^2-8x+7} dx$
 $= \int \sqrt{(x^2-8x+16)-9} dx$
 $= \int \sqrt{(x-4)^2-(3)^2} dx$

It is known that, $\int \sqrt{x^2-a^2} dx = \frac{x}{2}\sqrt{x^2-a^2} - \frac{a^2}{2} \log|x+\sqrt{x^2-a^2}| + C$

$\therefore I = \frac{(x-4)}{2}\sqrt{x^2-8x+7} - \frac{9}{2} \log|(x-4)+\sqrt{x^2-8x+7}| + C$

Hence, the correct Answer is D.

Exercise 7.8

Question 1:

$$\int_a^b x \, dx$$

Answer

It is known that,

$$\int_a^b f(x) \, dx = (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} [f(a) + f(a+h) + \dots + f(a+(n-1)h)], \text{ where } h = \frac{b-a}{n}$$

Here, $a = a$, $b = b$, and $f(x) = x$

$$\begin{aligned} \therefore \int_a^b x \, dx &= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} [a + (a+h) + \dots + (a+(n-1)h)] \\ &= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \left[\underbrace{(a+a+\dots+a)}_{n \text{ times}} + (h+2h+3h+\dots+(n-1)h) \right] \\ &= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} [na + h(1+2+3+\dots+(n-1))] \\ &= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \left[na + h \left\{ \frac{(n-1)(n)}{2} \right\} \right] \\ &= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \left[na + \frac{n(n-1)h}{2} \right] \\ &= (b-a) \lim_{n \rightarrow \infty} \frac{n}{n} \left[a + \frac{(n-1)h}{2} \right] \\ &= (b-a) \lim_{n \rightarrow \infty} \left[a + \frac{(n-1)h}{2} \right] \\ &= (b-a) \lim_{n \rightarrow \infty} \left[a + \frac{(n-1)(b-a)}{2n} \right] \\ &= (b-a) \lim_{n \rightarrow \infty} \left[a + \frac{\left(1 - \frac{1}{n}\right)(b-a)}{2} \right] \\ &= (b-a) \left[a + \frac{(b-a)}{2} \right] \\ &= (b-a) \left[\frac{2a+b-a}{2} \right] \\ &= \frac{(b-a)(b+a)}{2} \\ &= \frac{1}{2} (b^2 - a^2) \end{aligned}$$

Question 2:

$$\int_0^5 (x+1) dx$$

Answer

$$\text{Let } I = \int_0^5 (x+1) dx$$

It is known that,

$$\int_a^b f(x) dx = (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \left[f(a) + f(a+h) + \dots + f(a+(n-1)h) \right], \text{ where } h = \frac{b-a}{n}$$

Here, $a = 0$, $b = 5$, and $f(x) = (x+1)$

$$\Rightarrow h = \frac{5-0}{n} = \frac{5}{n}$$

$$\begin{aligned} \therefore \int_0^5 (x+1) dx &= (5-0) \lim_{n \rightarrow \infty} \frac{1}{n} \left[f(0) + f\left(\frac{5}{n}\right) + \dots + f\left((n-1)\frac{5}{n}\right) \right] \\ &= 5 \lim_{n \rightarrow \infty} \frac{1}{n} \left[1 + \left(\frac{5}{n} + 1\right) + \dots + \left\{ 1 + \left(\frac{5(n-1)}{n}\right) \right\} \right] \\ &= 5 \lim_{n \rightarrow \infty} \frac{1}{n} \left[\left(1 + 1 + 1 \dots 1\right) + \left[\frac{5}{n} + 2 \cdot \frac{5}{n} + 3 \cdot \frac{5}{n} + \dots + (n-1) \frac{5}{n}\right] \right] \\ &= 5 \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{5}{n} \{1 + 2 + 3 \dots (n-1)\} \right] \\ &= 5 \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{5}{n} \cdot \frac{(n-1)n}{2} \right] \\ &= 5 \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{5(n-1)}{2} \right] \\ &= 5 \lim_{n \rightarrow \infty} \left[1 + \frac{5}{2} \left(1 - \frac{1}{n}\right) \right] \\ &= 5 \left[1 + \frac{5}{2} \right] \\ &= 5 \left[\frac{7}{2} \right] \\ &= \frac{35}{2} \end{aligned}$$

Question 3:

$$\int_2^3 x^2 dx$$

Answer

It is known that,

$$\int_a^b f(x) dx = (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} [f(a) + f(a+h) + f(a+2h) \dots f\{a+(n-1)h\}], \text{ where } h = \frac{b-a}{n}$$

Here, $a = 2$, $b = 3$, and $f(x) = x^2$

$$\Rightarrow h = \frac{3-2}{n} = \frac{1}{n}$$

$$\begin{aligned} \therefore \int_2^3 x^2 dx &= (3-2) \lim_{n \rightarrow \infty} \frac{1}{n} \left[f(2) + f\left(2 + \frac{1}{n}\right) + f\left(2 + \frac{2}{n}\right) \dots f\left\{2 + (n-1)\frac{1}{n}\right\} \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \left[(2)^2 + \left(2 + \frac{1}{n}\right)^2 + \left(2 + \frac{2}{n}\right)^2 + \dots \left(2 + \frac{(n-1)}{n}\right)^2 \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \left[2^2 + \left\{ 2^2 + \left(\frac{1}{n}\right)^2 + 2 \cdot 2 \cdot \frac{1}{n} \right\} + \dots + \left\{ (2)^2 + \frac{(n-1)^2}{n^2} + 2 \cdot 2 \cdot \frac{(n-1)}{n} \right\} \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \left[\left(2^2 + \dots + 2^2 \right) + \left\{ \left(\frac{1}{n}\right)^2 + \left(\frac{2}{n}\right)^2 + \dots + \left(\frac{n-1}{n}\right)^2 \right\} + 2 \cdot 2 \cdot \left\{ \frac{1}{n} + \frac{2}{n} + \frac{3}{n} + \dots + \frac{(n-1)}{n} \right\} \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \left[4n + \frac{1}{n^2} \{1^2 + 2^2 + 3^2 \dots + (n-1)^2\} + \frac{4}{n} \{1 + 2 + \dots + (n-1)\} \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \left[4n + \frac{1}{n^2} \left\{ \frac{n(n-1)(2n-1)}{6} \right\} + \frac{4}{n} \left\{ \frac{n(n-1)}{2} \right\} \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \left[4n + \frac{n \left(1 - \frac{1}{n}\right) \left(2 - \frac{1}{n}\right)}{6} + \frac{4n-4}{2} \right] \\ &= \lim_{n \rightarrow \infty} \left[4 + \frac{1}{6} \left(1 - \frac{1}{n}\right) \left(2 - \frac{1}{n}\right) + 2 - \frac{2}{n} \right] \\ &= 4 + \frac{2}{6} + 2 \\ &= \frac{19}{3} \end{aligned}$$

Question 4:

$$\int_1^4 (x^2 - x) dx$$

Answer

$$\begin{aligned} \text{Let } I &= \int_1^4 (x^2 - x) dx \\ &= \int_1^4 x^2 dx - \int_1^4 x dx \end{aligned}$$

$$\text{Let } I = I_1 - I_2, \text{ where } I_1 = \int_1^4 x^2 dx \text{ and } I_2 = \int_1^4 x dx \quad \dots(1)$$

It is known that,

$$\int_a^b f(x) dx = (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} [f(a) + f(a+h) + f(a+(n-1)h)], \text{ where } h = \frac{b-a}{n}$$

$$\text{For } I_1 = \int_1^4 x^2 dx,$$

$$a = 1, b = 4, \text{ and } f(x) = x^2$$

$$\therefore h = \frac{4-1}{n} = \frac{3}{n}$$

$$\begin{aligned} I_1 &= \int_1^4 x^2 dx = (4-1) \lim_{n \rightarrow \infty} \frac{1}{n} [f(1) + f(1+h) + \dots + f(1+(n-1)h)] \\ &= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[1^2 + \left(1 + \frac{3}{n}\right)^2 + \left(1 + 2 \cdot \frac{3}{n}\right)^2 + \dots + \left(1 + \frac{(n-1)3}{n}\right)^2 \right] \\ &= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[1^2 + \left\{ 1^2 + \left(\frac{3}{n}\right)^2 + 2 \cdot \frac{3}{n} \right\} + \dots + \left\{ 1^2 + \left(\frac{(n-1)3}{n}\right)^2 + \frac{2 \cdot (n-1) \cdot 3}{n} \right\} \right] \\ &= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[\underbrace{\left(1^2 + \dots + 1^2\right)}_{n \text{ times}} + \left(\frac{3}{n}\right)^2 \{1^2 + 2^2 + \dots + (n-1)^2\} + 2 \cdot \frac{3}{n} \{1 + 2 + \dots + (n-1)\} \right] \end{aligned}$$

$$\begin{aligned}
 &= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{9}{n^2} \left\{ \frac{(n-1)(n)(2n-1)}{6} \right\} + \frac{6}{n} \left\{ \frac{(n-1)(n)}{2} \right\} \right] \\
 &= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{9n}{6} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + \frac{6n-6}{2} \right] \\
 &= 3 \lim_{n \rightarrow \infty} \left[1 + \frac{9}{6} \left(1 - \frac{1}{n} \right) \left(2 - \frac{1}{n} \right) + 3 - \frac{3}{n} \right] \\
 &= 3[1+3+3] \\
 &= 3[7]
 \end{aligned}$$

$$I_1 = 21 \quad \dots(2)$$

For $I_2 = \int_1^4 x dx$,

$a=1, b=4$, and $f(x) = x$

$$\Rightarrow h = \frac{4-1}{n} = \frac{3}{n}$$

$$\therefore I_2 = (4-1) \lim_{n \rightarrow \infty} \frac{1}{n} [f(1) + f(1+h) + \dots + f(a+(n-1)h)]$$

$$= 3 \lim_{n \rightarrow \infty} \frac{1}{n} [1 + (1+h) + \dots + (1+(n-1)h)]$$

$$= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[1 + \left(1 + \frac{3}{n} \right) + \dots + \left\{ 1 + (n-1) \frac{3}{n} \right\} \right]$$

$$= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[\left(\underset{n \text{ times}}{1+1+\dots+1} \right) + \frac{3}{n} (1+2+\dots+(n-1)) \right]$$

$$= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[n + \frac{3}{n} \left\{ \frac{(n-1)n}{2} \right\} \right]$$

$$= 3 \lim_{n \rightarrow \infty} \frac{1}{n} \left[1 + \frac{3}{2} \left(1 - \frac{1}{n} \right) \right]$$

$$= 3 \left[1 + \frac{3}{2} \right]$$

$$= 3 \left[\frac{5}{2} \right]$$

$$I_2 = \frac{15}{2} \quad \dots(3)$$

From equations (2) and (3), we obtain

$$I = I_1 + I_2 = 21 - \frac{15}{2} = \frac{27}{2}$$

Question 5:

$$\int_{-1}^1 e^x dx$$

Answer

$$\text{Let } I = \int_{-1}^1 e^x dx \quad \dots(1)$$

It is known that,

$$\int_a^b f(x) dx = (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} [f(a) + f(a+h) \dots f(a+(n-1)h)], \text{ where } h = \frac{b-a}{n}$$

Here, $a = -1$, $b = 1$, and $f(x) = e^x$

$$\therefore h = \frac{1+1}{n} = \frac{2}{n}$$



$$\begin{aligned}
 \therefore I &= (1+1) \lim_{n \rightarrow \infty} \frac{1}{n} \left[f(-1) + f\left(-1 + \frac{2}{n}\right) + f\left(-1 + 2 \cdot \frac{2}{n}\right) + \dots + f\left(-1 + \frac{(n-1)2}{n}\right) \right] \\
 &= 2 \lim_{n \rightarrow \infty} \frac{1}{n} \left[e^{-1} + e^{\left(-1 + \frac{2}{n}\right)} + e^{\left(-1 + 2 \cdot \frac{2}{n}\right)} + \dots + e^{\left(-1 + \frac{(n-1)2}{n}\right)} \right] \\
 &= 2 \lim_{n \rightarrow \infty} \frac{1}{n} \left[e^{-1} \left\{ 1 + e^{\frac{2}{n}} + e^{\frac{4}{n}} + e^{\frac{6}{n}} + e^{\frac{(n-1)2}{n}} \right\} \right] \\
 &= 2 \lim_{n \rightarrow \infty} \frac{e^{-1}}{n} \left[\frac{e^{\frac{2n}{n}} - 1}{e^{\frac{2}{n}} - 1} \right] \\
 &= e^{-1} \times 2 \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{e^2 - 1}{e^{\frac{2}{n}} - 1} \right] \\
 &= \frac{e^{-1} \times 2(e^2 - 1)}{\lim_{\frac{2}{n} \rightarrow 0} \left(\frac{e^{\frac{2}{n}} - 1}{\frac{2}{n}} \right) \times 2} \\
 &= e^{-1} \left[\frac{2(e^2 - 1)}{2} \right] \left[\lim_{h \rightarrow 0} \left(\frac{e^h - 1}{h} \right) = 1 \right] \\
 &= \frac{e^2 - 1}{e} \\
 &= \left(e - \frac{1}{e} \right)
 \end{aligned}$$

Question 6:

$$\int_0^4 (x + e^{2x}) dx$$

Answer

It is known that,

$$\int_a^b f(x) dx = (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \left[f(a) + f(a+h) + \dots + f(a+(n-1)h) \right], \text{ where } h = \frac{b-a}{n}$$

Here, $a = 0$, $b = 4$, and $f(x) = x + e^{2x}$

$$\therefore h = \frac{4-0}{n} = \frac{4}{n}$$

$$\begin{aligned}
 \Rightarrow \int_0^4 (x + e^{2x}) dx &= (4-0) \lim_{n \rightarrow \infty} \frac{1}{n} [f(0) + f(h) + f(2h) + \dots + f((n-1)h)] \\
 &= 4 \lim_{n \rightarrow \infty} \frac{1}{n} [(0 + e^0) + (h + e^{2h}) + (2h + e^{2 \cdot 2h}) + \dots + \{(n-1)h + e^{2(n-1)h}\}] \\
 &= 4 \lim_{n \rightarrow \infty} \frac{1}{n} [1 + (h + e^{2h}) + (2h + e^{4h}) + \dots + \{(n-1)h + e^{2(n-1)h}\}] \\
 &= 4 \lim_{n \rightarrow \infty} \frac{1}{n} [\{h + 2h + 3h + \dots + (n-1)h\} + \{1 + e^{2h} + e^{4h} + \dots + e^{2(n-1)h}\}] \\
 &= 4 \lim_{n \rightarrow \infty} \frac{1}{n} \left[h \{1 + 2 + \dots + (n-1)\} + \left(\frac{e^{2hn} - 1}{e^{2h} - 1} \right) \right] \\
 &= 4 \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{h(n-1)n}{2} + \left(\frac{e^{2hn} - 1}{e^{2h} - 1} \right) \right] \\
 &= 4 \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{4}{n} \cdot \frac{(n-1)n}{2} + \left(\frac{e^8 - 1}{e^2 - 1} \right) \right] \\
 &= 4(2) + 4 \lim_{n \rightarrow \infty} \left(\frac{\frac{e^8 - 1}{e^2 - 1}}{\frac{4}{n}} \right) \cdot 8 \\
 &= 8 + \frac{4 \cdot (e^8 - 1)}{8} \quad \left(\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1 \right) \\
 &= 8 + \frac{e^8 - 1}{2} \\
 &= \frac{15 + e^8}{2}
 \end{aligned}$$

